

Translation planes admitting a linear Abelian group of order $(q + 1)^2$.

Esteban Diazⁱ

*Department of Mathematics,
University of Connecticut.
diaz@math.uconn.edu*

Oscar Vega

*Department of Mathematics,
California State University, Fresno.
ovega@csufresno.edu*

Abstract. Translation planes of order q^2 and spread in $PG(3, q)$, where q is an odd prime power and $q^2 - 1$ has a p -primitive divisor, that admit a linear Abelian group of order $(q + 1)^2$ containing at most three kernel homologies are shown to be associated to flocks of quadratic cones.

Keywords: Translation plane, flock of quadratic cone, homologies

MSC 2000 classification: Primary 51, 05; Secondary 20

1 Introduction

In a series of papers, that span more than fifteen years, translation planes of order q^2 with spread in $PG(3, q)$ that admit a linear collineation group G of order $q(q + 1)$ were completely classified as associated to conical flocks planes. It is also known that translation planes of order q^2 , with spread in $PG(3, q)$, that admit a cyclic homology group of order $q + 1$ are equivalent to conical flocks planes. Moreover, in this situation is possible to show that the full collineation group of the translation plane that admit the cyclic homology of order $q + 1$ also admits a group of order $(q + 1)^2$. In the spirit of this idea, it is possible to argue that the translation planes that admit a group of order $(q + 1)^2$ may also be associated with conical flocks planes.

We will use standard notation and results found in the literature on finite translation planes and/or flocks of quadratic cones. More details may be found in [2, 10, 11]. In particular, we will use André's [1] theory of translation planes and spreads of vector spaces. A collineation of a translation plane π is a one-to-one mapping of the points onto the points of π that preserves incidence. The collineation group of π is denoted $\text{Aut}(\pi)$, and the stabilizer of 0 is called the

ⁱThis article is dedicated to Norman Johnson on the occasion of his 70th birthday.

translation complement of π . If $\Psi \in \text{Aut}(\pi)$ fixes a line l pointwise and all the lines through a point P setwise, then Ψ is called a perspectivity of π , if $P \in l$ then Ψ is called an elation, otherwise it is called a homology. In either case, P is called the center of Ψ and l is called the axis of Ψ .

1 Theorem. [Johnson, [6]] *Translation planes with spreads in $PG(3, q)$ admitting cyclic affine homology groups of order $q + 1$ are equivalent to flocks of quadratic cones.*

2 Theorem. [Johnson [7]] *Let V be a vector space of dimension $2n$ over a field $F \cong GF(q)$, for $q = p^r$, p a prime. Assume that a collineation $\sigma \in GL(2n, q)$ has order dividing $q^n - 1$ but not dividing $q^t - 1$ for $t < n$. If σ fixes at least three mutually disjoint n -dimensional F -subspaces then there is an associated Desarguesian spread Σ admitting σ as a kernel homology. Furthermore, the normalizer of $\langle \sigma \rangle$ is a collineation group of Σ . Let us call Σ an ‘Ostrom phantom’.*

The problem we will study in this paper is:

3 Problem. [Il problema Abeliano rosso] Determine the translation planes π of order q^2 with spread in $PG(3, q)$ that admit an Abelian collineation group G of order $(q + 1)^2$ in $GL(4, q)$.

A conjecture regarding this problem says that planes such as those described above must be associated to a flock of a quadratic cone. This comes after a series of papers (see for example [3–5, 8, 9]) that completely classified translation planes of order q^2 with spread in $PG(3, q)$ admitting a collineation group $G \subset GL(4, q)$ of order $q(q + 1)$. Such translation planes turned out to be conical flock planes or derived conical flock planes, except in a few sporadic cases, see [3] for more details. Also, it was shown that the group G is solvable and that it has a subgroup H of order $q + 1$ that normalizes an elation subgroup E of order q . Moreover, when G fixes two components of π there is an Ostrom phantom Σ associated to π , G is in $GL(2, q^2)$, and H fixes at least two components of π (one being the elation axis). It follows that H and G fix a regulus of the flock’s plane.

Theorem 1 implies that if a translation plane Π with spread in $PG(3, q)$ admits a regulus inducing affine homology H_1 of order $q + 1$ in the translation complement, for example H_1 cyclic, then Π is equivalent to a conical flock plane \mathcal{F} . Now if one takes the normalizer N of H_1 , then the quotient group N/H_1 acts as a collineation group of \mathcal{F} , permuting q reguli and fixing one of the reguli of \mathcal{F} . Connecting the previous paragraph with this idea we would have that N/H_1 has a subgroup of order $(q + 1)$. Hence, the normalizer N should contain a subgroup of order $(q + 1)^2$. This justifies the conjecture.

Our main result follows, it will be proved as a series of results in the next

section.

4 Theorem. *Let π be a translation plane of order q^2 (q an odd prime power) with spread in $PG(3, q)$ admitting a linear Abelian collineation group G of order $(q+1)^2$. Assume that G contains at most three kernel homologies and that $q^2 - 1$ admits a p -primitive divisor, then π is associated to a conical flock plane.*

5 Remark. Johnson and Pomareda [8] prove that under the same condition of theorem 4, in the case q even, the translation plane admitting the collineation group of order $(q+1)^2$ is André or Desarguesian.

2 Proof of the main theorem

For the rest of this article we will assume the hypothesis of theorem 4. Also, S is a spread of π and u is a p -primitive divisor of $q^2 - 1$.

6 Lemma. *Any Sylow u -subgroup S_u of G fixes 2 components of π .*

PROOF. Note that $u \neq 2$. Now let u^{2a} be the maximal power of u dividing $(q+1)^2$.

Since $q^2 + 1 = (q+1)(q-1) + 2$, then $(q^2 + 1, u^{2a}) = 1$. It follows that the action of S_u on the components of S must fix at least one component.

Now S_u acts on q^2 components of S , but since $(q^2, u^{2a}) = 1$ then S_u must fix a second component. ◻

7 Lemma. *Suppose an element $g \in S_u$ fixes a non-zero point in a component L that is being fixed by S_u . Then g is an affine homology with axis L .*

PROOF. Since g is linear, under the hypothesis given we have that g must fix a 1-dimensional $GF(q)$ -subspace A of L . Now using that $(q, u^{2a}) = 1$ we get that A has a 1-dimensional Maschke complement B .

Now recall that the order of g is a power of u , and that the number of non-zero elements in A (and B) is $q - 1$. So, since $(q - 1, u^t) = 1$ for any integer t , we have that g must fix a point in A (and B), and thus g must fix A and B pointwise. Hence, g fixes the component L pointwise. ◻

Now we change basis, if necessary, to get the two components that are fixed by S_u to be $x = 0$ and $y = 0$. Then we consider S_u acting on the 1-dimensional subspaces of $x = 0$. Since the order of S_u is u^{2a} , and it is acting on a set with $q + 1 = u^a r$ elements, where $(r, u) = 1$ and $u^{2a} > u^a$, then the stabilizer of at least one of these 1-dimensional subspaces must be non-trivial. Using the ‘Maschke argument’ used in the proof of the previous lemma we can assure that there is a subgroup of S_u fixing $x = 0$ pointwise, call $H_{x=0}^{(u)}$ to be the largest such a subgroup. Similarly, $H_{y=0}^{(u)}$ is the largest subgroup of S_u fixing every point in $y = 0$. Moreover, they are normal in S_u and, by lemma 7, homology groups.

8 Lemma. $H_{x=0}^{(u)}$ and $H_{y=0}^{(u)}$ are cyclic. $S_u = H_{x=0}^{(u)} \oplus H_{y=0}^{(u)}$.

PROOF. Since homology groups are Frobenius complements (see [12], for example), and Frobenius complements have cyclic odd-order Sylow subgroups, then both $H_{x=0}^{(u)}$ and $H_{y=0}^{(u)}$ are cyclic.

If we look at the orbit equation of the action of S_u on the 1-dimensional subspaces of $x = 0$ (let's call them p_i 's) we get

$$u^a r = q + 1 = \sum \frac{u^{2a}}{|Stab(p_i)|}$$

where the sum considers only one p_i per orbit under S_u and $(r, u) = 1$. We notice that none of the summands can equal one because S_u cannot contain nontrivial elements that are homologies with two different axes. Also, if all the stabilizers contain less than u^a elements, then $(r, u) \neq 1$. It follows that at least one of the stabilizers has at least u^a elements. Since any element fixing a 1-dimensional subspace of $x = 0$ fixes $x = 0$ pointwise, then all stabilizers have at least u^a elements. It follows that $|H_{x=0}^{(u)}| \geq u^a$ and, similarly, $|H_{y=0}^{(u)}| \geq u^a$. Hence, $|H_{x=0}^{(u)} \cap H_{y=0}^{(u)}| = 1$ implies $S_u = H_{x=0}^{(u)} \oplus H_{y=0}^{(u)}$. □ QED

9 Remark. Note that $H_{x=0}^{(u)}$ and $H_{y=0}^{(u)}$ commuting implies that they are symmetric homology groups.

Also, the previous three lemmas are valid even when G is not Abelian.

10 Theorem. *There is $g \in S_u$ of order u that fixes 3 components of S (two of them being $x = 0$ and $y = 0$). Furthermore, there is an Ostrom phantom Σ induced by g , and $G \leq \Gamma L(2, q^2)$.*

PROOF. We know $H_{x=0}^{(u)}$ is an affine homology group with axis $x = 0$ and coaxis $y = 0$ that acts on the remaining $q^2 - 1$ components of the given spread producing $(q^2 - 1)/u^a$ orbits. Note that $H_{y=0}^{(u)}$ acts on these orbits and that, since $(u^a, (q^2 - 1)/u^a) = 1$ then $H_{y=0}^{(u)}$ fixes at least one of them, call it M . Then, we can consider S_u acting on M . The orbit equation of this action is:

$$u^a = \sum \frac{u^{2a}}{|Stab(l_i)|}$$

where the sum is on the components of M , one l_i per orbit under S_u .

It is clear that none of the stabilizers can be trivial. In this way we obtain an element $g \in S_u$ of order u that fixes some l_i , $x = 0$ and $y = 0$. This element g satisfies the hypothesis of theorem 2, and thus there is an Ostrom phantom Σ and the normalizer of $\langle g \rangle$ in $GL(4, q)$ is a collineation group of Σ . Since G is Abelian then it is a collineation group of Σ . □ QED

11 Lemma. G has a subgroup of order $(q+1)^2/4$ which is the direct sum of two cyclic symmetric affine homology groups of order $(q+1)/2$. Their axes/coaxes are $x=0$ and $y=0$.

PROOF. Let us denote by $(q+1)_2$ the maximal power of 2 in $q+1$, and by S_2 the 2-Sylow subgroup of G , which has order $(q+1)_2$. Since G is Abelian, then S_2 acts on the fixed points of $H_{x=0}^{(u)}$, which forces S_2 to fix $x=0$. Similarly, S_2 fixes $y=0$.

Now, consider the restriction of S_2 to $x=0$, this group is isomorphic to $S_2/(S_2)_{x=0}$, where $(S_2)_{x=0}$ is the subgroup of S_2 that fixes $x=0$ pointwise. But, also $S_2/(S_2)_{x=0}$ is a subgroup of $PGL(2, q)$, since it acts on $y=0$, which can be regarded as a Desarguesian plane. It follows that

$$\left| \frac{S_2}{(S_2)_{x=0}} \right| = \frac{(q+1)_2^2}{|(S_2)_{x=0}|}$$

divides

$$|PGL(2, q)|_2 = (q-1)_2(q+1)_2 = 2(q+1)_2.$$

From that we get

$$\frac{(q+1)_2}{2} \mid |(S_2)_{x=0}|.$$

Now consider $H < G$ of order $(q+1)_{odd}^2 = (q+1)^2/(q+1)_2$. Using the same arguments used with $(S_2)_{x=0}$ we obtain

$$(q+1)_{odd} \mid |H_{x=0}|.$$

Hence, the subgroup $G_{x=0}$ of G that fixes $x=0$ pointwise has order a multiple of

$$|(S_2)_{x=0}| |H_{x=0}| = \frac{(q+1)_2}{2} (q+1)_{odd} = \frac{(q+1)}{2}$$

Using the same argument with $y=0$ we obtain that $G_{x=0} \times G_{y=0}$ is the desired subgroup of G . □

12 Corollary. Let $G_{x=0}$ and $G_{y=0}$ be the cyclic homology groups of order $(q+1)/2$ found in lemma 11. Then, after a change of basis if necessary, $G = G_1 \times G_2$ with

- (1) $G_{x=0} < G_1 \cong \mathbb{Z}_{q+1}$ and $G_{y=0} < G_2 \cong \mathbb{Z}_{q+1}$,
- (2) $G_{x=0} < G_1 \cong \mathbb{Z}_{2(q+1)}$, $G_{y=0} = G_2 \cong \mathbb{Z}_{\frac{q+1}{2}}$, and $(q+1)/2$ is odd, or
- (3) $G \cong \mathbb{Z}_{2(q+1)} \times \mathbb{Z}_{\frac{q+1}{2}}$, $(q+1)/2$ is even, and at least one of $G_{x=0}$ or $G_{y=0}$ is a subgroup of one factor of G .

PROOF. If $(q+1)/2$ is odd, then the 2-Sylow subgroup of G intersects $G_{x=0} \times G_{y=0}$ trivially, and thus we are done.

If $(q+1)/2$ is even then having G to be the direct product of three or more factors would force Σ to admit an elementary Abelian group of order 8 or 16, this is a contradiction. It follows that either $G \cong \mathbb{Z}_{q+1} \times \mathbb{Z}_{q+1}$ or $G \cong \mathbb{Z}_{2(q+1)} \times \mathbb{Z}_{\frac{q+1}{2}}$.

In the former case we break $G_{x=0} \times G_{y=0}$ into the direct product of its 2-Sylow subgroup H_2 with its complement. The idea used in the case $(q+1)/2$ odd is applicable to the complement of the H_2 , thus we will look at this group.

Assume that $H_2 \cong \mathbb{Z}_{2^n} \times \mathbb{Z}_{2^n}$ is a subgroup of $G_2 \cong \mathbb{Z}_{2^{n+1}} \times \mathbb{Z}_{2^{n+1}}$ (the 2-Sylow subgroup of G). An element in G_2 of order 2^n looks like (\bar{a}, \bar{b}) where both a and b are congruent to 2 modulo 4, it follows that $(\bar{a}, \bar{b}) = 2(\bar{a}/2, \bar{b}/2)$. Hence, each of the \mathbb{Z}_{2^n} 's in H_2 is contained in some $\mathbb{Z}_{2^{n+1}}$ in G_2 , thus a change of generators of G_2 implies that we can consider each of the factors of H_2 to be contained in one of the factors of G_2 , which is what we wanted.

If $G \cong \mathbb{Z}_{2(q+1)} \times \mathbb{Z}_{\frac{q+1}{2}}$ and $(q+1)/2$ we break again $G_{x=0} \times G_{y=0}$ into the direct product of its 2-Sylow subgroup H_2 with its complement. In this case, an element of order 2^n can be either four times an element in G , twice but not four times an element in G , or not twice an element in G . If a generator of one of the factors of H_2 is either the first or third case, then we assure that one of the factors of $G_{x=0} \times G_{y=0}$ is a subgroup of one of the factors of G .

Let us assume that both of the generators of the factors of H_2 are twice but not four times an element of G . These elements look like $(\overline{8a+4}, \overline{4b+2})$ and $(\overline{8\alpha+4}, \overline{4\beta+2})$, where $a, b, \alpha, \beta \in \mathbb{Z}$. However,

$$2^{n-1}(\overline{8a+4}, \overline{4b+2}) = 2^{n-1}(\overline{8\alpha+4}, \overline{4\beta+2}) = (\overline{2^{n+1}}, \bar{0})$$

and thus the groups these elements generate intersect non-trivially. That is a contradiction. \square

13 Lemma. *The elements in G have the form*

$$(x, y) \rightarrow (xa, yb)$$

where $a, b \in GF(q^2)$. In particular, $G < GL(2, q^2)$.

PROOF. We know there exists an Ostrom phantom Σ associated to π , and that G is a collineation group of Σ that fixes $x = 0$ and $y = 0$. Then, a generic element $g \in G$ looks like

$$g : (x, y) \rightarrow (x^\sigma a, y^\sigma b)$$

where σ is 1 or q , and $a, b \in GF(q^2)^*$.

We also know that $H_{y=0}^{(u)}$ is a cyclic homology group of π , thus it is generated by

$$\tau : (x, y) \rightarrow (x, yc)$$

where $|c|$ must divide $(q+1)_{\text{odd}}$, which forces $c^\sigma = c$. Using that G is Abelian we check $g^{-1}\tau^{-1}g\tau$ for $g \in G$, as above, to obtain that $\sigma = 1$ for all $g \in G$. \square

14 Theorem. $G_{x=0} < G_1$ and $G_{y=0} < G_2$.

PROOF. We just need to show that in the case $G_1 \cong \mathbb{Z}_{2(q+1)}$, $G_2 \cong \mathbb{Z}_{(q+1)/2}$, and $(q+1)/2$ even, it is true that $G_{x=0} \subset G_1$ if and only if $G_{y=0} = G_2$.

Assume that $G_{x=0} \subset G_1 \cong \mathbb{Z}_{2(q+1)}$. Using the previous lemma we can assume that $G_1 = \langle f : (x, y) \mapsto (ax, by) \rangle$ and $G_{x=0} = \langle f^4 : (x, y) \mapsto (a^4x, y) \rangle$, where $|a| = 2(q+1)$, and $|b|$ is divisible by 4. Similarly, $G_2 = \langle g : (x, y) \mapsto (cx, dy) \rangle$ where $\text{lcm}(|c|, |d|) = (q+1)/2$.

Recall that $a, b, c, d \in GF(q^2)$ and note that that a^4 has order $(q+1)/2$, thus $c = a^{4i}$ for some positive integer i . So, if the order of d were a proper divisor of $(q+1)/2$ then

$$e \neq g^{|d|} : (x, y) \mapsto (c^{|d|}, y)$$

which is $f^{4|d|} \in G_1$, that is a contradiction. Hence, $|d| = (q+1)/2$, and thus the element $f^{-4i}g$ generates $G_{x=0}$.

Now assume that $G_{y=0} = G_2 \cong \mathbb{Z}_{(q+1)/2}$. Using the previous lemma we can assume that $G_2 = \langle g : (x, y) \mapsto (x, by) \rangle$, where $|b| = (q+1)/2$. We can also say that G_1 is generated by $f : (x, y) \mapsto (\alpha x, \beta y)$, where $\alpha, \beta \in GF(q^2)$ and $\text{lcm}(|\alpha|, |\beta|) = 2(q+1)$.

Note that $f^4 : (x, y) \mapsto (\alpha^4x, \beta^4y)$ has order $(q+1)/2$, and thus both α^4 and β^4 are powers of b . Hence, composing f^4 with some power of g gives us the collineation of G

$$(x, y) \mapsto (\alpha^4x, y)$$

which is a homology of order $(q+1)/2$. It follows that G is spanned by g and $h : (x, y) \mapsto (ax, cy)$, where the order of c is divisible by 4. It is clear that $G_{y=0}$ is contained in $\langle h \rangle$, which will be our new G_1 . \square

We now investigate each of the cases described in corollary 12 separately. Our goal is to show that π admits a cyclic affine homology group of order $q+1$, as having this will force π to be associated to a conical flock by [6].

When $G_{x=0} < G_1 \cong \mathbb{Z}_{q+1}$ and $G_{y=0} < G_2 \cong \mathbb{Z}_{q+1}$, we represent G as follows

$$G = \langle f : (x, y) \rightarrow (x\alpha, ya) \rangle \times \langle g : (x, y) \rightarrow (xb, y\beta) \rangle$$

where a and b have order $q+1$, and α and β have order 2. Also, without loss of generality, $G_{x=0} \subset \langle f \rangle$ and $G_{y=0} \subset \langle g \rangle$.

15 Lemma. G is the direct sum of two symmetric cyclic affine homology groups of order $q + 1$.

PROOF. If $(q + 1)/2$ is even. Note that $f^{-(q+1)/2}g$ is defined by $(x, y) \rightarrow (xb, y)$, which is a homology of order $q + 1$. Similarly, $g^{-(q+1)/2}f$ is a homology of order $q + 1$ that is symmetric to $f^{-(q+1)/2}g$.

If $(q + 1)/2$ is odd. Compose f and g with $(x, y) \rightarrow (-x, -y)$ to obtain

$$(x, y) \rightarrow (x, -ya) \quad \text{and} \quad (x, y) \rightarrow (-xb, y)$$

It is clear that these elements are affine homologies of order $q + 1$ that generate G and that have symmetric axis/coaxis. \square

Now we will look at $G \cong \mathbb{Z}_{2(q+1)} \times \mathbb{Z}_{(q+1)/2}$. In this case

$$\mathbb{Z}_{2(q+1)} = \langle f : (x, y) \mapsto (x\alpha, ya) \rangle \quad \mathbb{Z}_{(q+1)/2} = \langle g : (x, y) \rightarrow (xb, y) \rangle$$

where $a, b, \alpha \in GF(q^2)$ with $\text{lcm}[|a|, |\alpha|] = 2(q + 1)$, $|b| = (q + 1)/2$ and, since $f^4 \in G_{y=0}$, $|\alpha|$ divides 4.

16 Theorem. G admits a cyclic homology group of order $q + 1$.

PROOF. If $|\alpha| = 1$ or 2 then $f^2 \in G_{y=0}$ and thus the plane admits a cyclic homology group of order $q + 1$.

If $|\alpha| = 4$ and $(q + 1)/2$ is even, then $\alpha^2 = -1 = b^{(q+1)/4}$. It follows that $g^{(q+1)/4}f^2 \in G_{y=0}$ and it has order $q + 1$.

Now assume that $|\alpha| = 4$ and $(q + 1)/2$ is odd. In the case of $|a| = (q + 1)/2$, then $f^{(q+1)/2} \in G_{x=0}$. It follows that $f^{(q+1)/2}g \in G_{x=0}$ and has order $2(q + 1)$. Similarly, if $|a| = q + 1$, then $f^{q+1} \in G_{x=0}$, and thus $f^{q+1}g \in G_{x=0}$ and has order $q + 1$. In either case the plane admits a cyclic homology group of order $q + 1$.

Finally, consider $|\alpha| = 4$, $|a| = 2(q + 1)$, and $(q + 1)/2$ odd. Note that $|\langle \alpha b \rangle| = 2(q + 1)$. So, there is a positive integer i such that either $a = \alpha b^i$ or $a^{-1} = \alpha b^i$.

In the first case, $f^{(q+1)/2}$ is a kernel homology of order 4, which contradicts our hypothesis.

In the second case, $a^{-1} = \alpha b^i$ forces $a^{-(q+1)/2} = \alpha^{(q+1)/2} = \alpha^{\pm 1}$. We will now look at these two cases separately.

If $a^{-(q+1)/2} = \alpha$, we use $\alpha^{-1} = \alpha b^i$ to get $\alpha^{-(q+1)/2} = a^{(q+1)/2}$. It follows that $\alpha^{(q+1)/2} = \alpha$. All this implies that $f^{(q+1)/2}$ is defined by

$$f^{(q+1)/2} : (x, y) \mapsto (x\alpha, y\alpha^{-1})$$

Since $(q + 1)/2$ odd then $q \equiv 1 \pmod{3}$, thus $\alpha \in GF(q)$. It follows that the map

$$\sigma : (x, y) \mapsto (x\alpha, y\alpha)$$

is a kernel homology, which put together with $f^{(q+1)/2}$ yields

$$\sigma f^{(q+1)/2} : (x, y) \mapsto (-x, y)$$

Hence, the map $\sigma f^{(q+1)/2} g$ is an element of $G_{x=0}$ of order $q + 1$.

Finally, if $a^{(q+1)/2} = \alpha$, just as we did above, we get $\alpha^{(q+1)/2} = \alpha^{-1}$ and thus

$$f^{(q+1)/2} : (x, y) \mapsto (x\alpha^{-1}, y\alpha)$$

The kernel homology $\tau : (x, y) \mapsto (x\alpha^{-1}, y\alpha^{-1})$ finishes the proof. \square *QED*

17 Remark. The kernel homologies assumption of theorem 4, is minimal in the sense that, as the situation of two cyclic homology groups of order $q + 1$ shows, we have a kernel homology of order 2 in the group G . It is an open problem to determine what happens when four or more kernel homologies are in G .

References

- [1] J. ANDRÉ: *Über Perspektivitäten in endlichen affinen Ebenen (German)*, Arch. Math., **9** (1958), 228–235.
- [2] M. BILIOTTI, V. JHA, N. L. JOHNSON: *Handbook of Finite Translation Planes*, Pure and Applied Mathematics (Boca Raton), 289. Chapman & Hall/CRC, Boca Raton, FL, 2007.
- [3] V. JHA, N. L. JOHNSON: *The classification of spreads in $PG(3, q)$ admitting linear groups of order $q(q + 1)$ I: Odd order*, J. Geom., no. 1-2 **81** (2004), 46–80.
- [4] V. JHA, N. L. JOHNSON: *Multiple nests*, Note Mat., no. 1 **23** (2004-05), 123–149.
- [5] V. JHA, N. L. JOHNSON: *Nests of reguli and flocks of quadratic cones*, Simon Stevin, no. 3-4 **63** (1989), 311–338.
- [6] N.L. JOHNSON: *Homology groups of translation planes and flocks of quadratic cones, I: The structure*, Bull. Belg. Math. Soc. Simon Stevin, no. 5 **12** (2006), 827–844.
- [7] N.L. JOHNSON: *Translation planes of order q^2 that admit $q + 1$ elations*, Geom. Dedicata, no. 4 **15** (1984), 329–337.
- [8] N.L. JOHNSON, R. POMAREDA, *André planes and nests of reguli*, Geom. Dedicata, no. 3 **31** (1989), 245–260.
- [9] N. L. JOHNSON, R. POMAREDA: *Mixed Nests*, J. Geom., no. 1-2 **56** (1996), 59–86.
- [10] H. LÜNEBURG: *Translation Planes*, Springer-Verlag, Berlin–New York, 1980.
- [11] T. G. OSTROM: *Finite Translation Planes*, Lecture Notes in Mathematics, No. 158, Berlin-Heidelberg-New York, Springer-Verlag, 1970.
- [12] D.S. PASSMAN: *Permutation Groups*, Benjamin, New York, 1968.
- [13] J. ANDRÉ: *Über nicht-Desarguesian Ebenen mit transitiver translation gruppen*, Math. Z. **60**, 156–186.
- [14] R. D. BAKER, G. L. EBERT, T. PENTILIA: *Hyperbolic fibrations and q -clans*, Des. Codes Cryptogr., no. 2-3 **34** (2005), 295–305.

- [15] G. L. EBERT: *Spreads Admitting Regular Elliptic Covers*, Europ. J. Comb., no. 4 **10** (1989), 319–330.
- [16] G. L. EBERT: *Nests, Covers, and Translation Planes*, Proc. 11th British Combinatorial Conference (London, 1987), Ars Combin., no. C **25** (1988), 213–233.
- [17] N.L. JOHNSON: *Subplane Covered Nets*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 222, Marcel Dekker, Inc., New York, 2000.
- [18] K. ZSIGMONDY: ZUR THEORIE DER POTENZRESTE, Monatshefte Math. Phys., **3** (1982), 265–284.