

and

$$N_1^2 - (2q+2-g)N_1 + (q+1)^2 - (q^2+1)g - 2qg^2 \leq 0,$$

from which the result follows.

For  $g > \frac{1}{2}(q-\sqrt{q})$ , Ihara's result is better than Serre's.

#### 4. THE ESSENTIAL IDEA IN A PARTICULAR CASE

Let  $\mathcal{C}$  be as in §2, but consider it as a curve over  $\bar{K}$ , the algebraic closure of  $K = GF(q)$ . Also suppose that  $\mathcal{C}$  is embedded in the plane  $PG(2, \bar{K})$  and let  $\varphi$  be the Frobenius map given by

$$P(x_0, x_1, x_2)\varphi = P(x_0^q, x_1^q, x_2^q)$$

where  $P(x_0, x_1, x_2)$  is the point of the plane with coordinate vector  $(x_0, x_1, x_2)$ . Then

$$\begin{aligned} \mathcal{C} &= V(F) \\ &= \{P(x_0, x_1, x_2) \mid F(x_0, x_1, x_2) = 0\} \end{aligned}$$

for some form  $F$  in  $K[X_0, X_1, X_2]$ . Also  $\mathcal{C}\varphi = \mathcal{C}$  and the points of  $\mathcal{C}$  rational over  $GF(q)$  are exactly the fixed points of  $\varphi$  on  $\mathcal{C}$ .

For any non-singular point  $P = P(x_0, x_1, x_2)$  the tangent  $T_p$  at  $P$  is

$$T_p = V\left(\frac{\partial F}{\partial x_0} X_0 + \frac{\partial F}{\partial x_1} X_1 + \frac{\partial F}{\partial x_2} X_2\right).$$

In affine coordinates,

$$T_p = V\left(\frac{\partial f}{\partial a}(x-a) + \frac{\partial f}{\partial b}(x-b)\right)$$

where  $f(x,y) = F(x,y,1)$ .

Instead of looking at fixed points of  $\varphi$ , let us look at the set of points such that  $P\varphi \in T_p$ . As  $P \in T_p$ , this set contains the  $\text{GF}(q)$ -rational points of  $\mathcal{C}$ . Let

$$h = (x^q - x)f_x + (y^q - y)f_y.$$

Then

$$\begin{aligned} h_x &= (qx^{q-1} - 1)f_x + (x^q - x)f_{xx} + (y^q - y)f_{yx} \\ &= -f_x + (x^q - x)f_{xx} + (y^q - y)f_{yx} \end{aligned}$$

and

$$h_y = -f_y + (x^q - x)f_{xy} + (y^q - y)f_{yy}.$$

So  $V(h)$  and  $V(f)$  have a common tangent at any  $\text{GF}(q)$ -rational point of  $\mathcal{C}$  that is non-singular. So, if  $N$  is the number of  $\text{GF}(q)$ -rational points of  $\mathcal{C}$  and the degree of  $f$  is  $d$ , then Bézout's theorem implies, when  $f$  is not a component of  $h$ , that

$$\begin{aligned} (d+q-1)d &= \deg h \deg f \\ &= \text{sum of the intersection numbers at} \\ &\quad \text{the points of } V(f) \cap V(h) \\ &\geq 2N. \end{aligned}$$

Hence  $N \leq \frac{1}{2}d(d+q-1)$ .

Now, suppose that  $V(f)$  is a component of  $V(h)$ , or equivalently that  $h=0$  as a function on  $V(f)$ . Therefore

$$\begin{aligned} (x^q - x)f_x/f_y + (y^q - y) &= 0, \\ (x^q - x)\frac{dy}{dx} - (y^q - y) &= 0. \end{aligned}$$

Differentiating gives

$$(x^q - x) \frac{d^2 y}{dx^2} - \frac{dy}{dx} - \frac{d}{dx}(y^q - y) = 0$$

Remembering that  $\frac{d}{dx} = \frac{\partial}{\partial x} + \frac{dy}{dx} \frac{\partial}{\partial y}$ , we obtain that

$$(x^q - x) \frac{d^2 y}{dx^2} = 0$$

$$\frac{d^2 y}{dx^2} = 0.$$

Since  $\frac{dy}{dx} = -f_x/f_y$ , it follows that

$$\frac{d^2 y}{dx^2} = -f_y^{-2} \{f_{xx} f_y^2 - 2f_{xy} f_x f_y + f_{yy} f_x^2\}.$$

**THEOREM 4.1:** If  $\frac{d^2 y}{dx^2} \neq 0$ , that is,  $\mathcal{C}$  is not all inflexions and  $q$  is odd, then  $N \leq \frac{1}{2} d(d+q-1)$ .

In fact  $\frac{d^2 y}{dx^2} = 0$  can only occur when  $\mathcal{C}$  is a line or the characteristic  $p \leq d$ . For example, when  $f = x^{p^r+1} + y^{p^r+1}$ , then  $\mathcal{C}$  is all inflexions. A particular case of this phenomenon is the Hermitian curve  $\mathcal{U}_{2,q} = V(X_0^{\sqrt{q}+1} + X_1^{\sqrt{q}+1} + X_2^{\sqrt{q}+1})$  when  $q$  is a square.

Since every curve of genus 3 can be embedded in the plane as a non-singular quartic, we can see how theorem 4.1 compares with Serre's bound for  $N_q(3)$  and its actual value.

q	3	5	7	9	11	13	17	19
$2(q+3)$	12	16	20	24	28	32	40	44
$q+1+3[2\sqrt{q}]$	13	18	23	28	30	35	42	44
$N_q(3)$	10	16	20	28	28	32	40	44

Thus, for  $q$  odd with  $q \leq 19$  and  $q \neq 3$  or  $9$ , the theorem gives the best possible result. A curve achieving  $N_q(3)$  is  $\mathcal{W}_{2,9}$ .

### 5. WEIERSTRASS POINTS IN CHARACTERISTIC ZERO.

First consider the canonical curve  $\mathcal{C}^{2g-2}$  of genus  $g \geq 3$  in  $PG(g-1, \mathbb{C})$ . The Weierstrass points, W-points for short, are the points at which the osculating hyperplane has  $g$  coincident intersections. In this case, with  $w$  the number of W-points

$$w = g(g^2 - 1).$$

In any case,

$$2g + 2 \leq w \leq g(g^2 - 1)$$

with the lower bounded achieved only for hyperelliptic curves.

A curve of genus  $g > 1$  is hyperelliptic if it has a linear series  $\gamma_{\frac{1}{2}}$  (a 2-sheeted covering) on it; for example, a plane quartic with a double point. It has equation

$$y^2 = f(x)$$

with genus  $g = [\frac{1}{2}(d-1)]$  where  $d = \deg f$ .

Consider the case  $g=3$  of the canonical curve  $\mathcal{C}^4$ , a non-singular plane quartic. The W-points are the 24 inflexions. We note that